arXiv:1406.2359v1 [gr-qc] 4 Dec 2013

The Stability of a Shearing Viscous Star with Electromagnetic Field

M. Sharif[1] *and M. Azam[1,2] †
[1] Department of Mathematics, University of the Punjab,
Quaid-e-Azam Campus, Lahore-54590, Pakistan.
[2] Division of Science and Technology, University of Education,
Township Campus, Lahore-54590, Pakistan.

Abstract

We analyze the role of electromagnetic field for the stability of shearing viscous star with spherical symmetry. Matching conditions are given for the interior and exterior metrics. We use perturbation scheme to construct the collapse equation. The range of instability is explored in Newtonian and post Newtonian (pN) limits. We conclude that the electromagnetic field diminishes the effects of shearing viscosity in the instability range and makes the system more unstable at both Newtonian and post Newtonian approximations.

Keywords: Gravitational collapse; Electromagnetic field; Instability.
PACS: 04.20.-q; 04.25.Nx; 04.40.Dg; 04.40.Nr.

1 Introduction

Charged self-gravitating objects have the tendency of undergoing many phases during gravitational collapse resulting a charged black holes or naked singularities [1]. The stability of these exact solutions is an interesting subject under the perturbation scheme. It is believed that these stars with huge

*msharif.math@pu.edu.pk
†azammath@gmail.com

charge cannot be stable [2, 3]. However, the electric charge has great relevance during the structure formation and evolution of astrophysical objects [4]-[11].

A stellar model may be stable in one phase and later becomes unstable in another phase. The stability of this model is subjected against any disturbance. Dynamical instability of self-gravitating objects is interrelated with structure as well as evolution of astrophysical objects. In this scenario, Chandrasekhar [12] investigated the problem of dynamical instability for the isotropic perfect fluid of a pulsating system and found the instability range in terms of adiabatic index $\Gamma < \frac{4}{3}$.

It is well discussed in literature that the instability range of the fluid would be decreased or increased through different physical properties of the fluid. In this context, the dynamical instability for adiabatic, non-adiabatic, anisotropic and shearing viscous fluids have been explored [13]-[17]. Chan [18] found that both pressure anisotropy and effective adiabatic index are increased by the shearing viscosity in a collapsing radiating star. Horvat et al. [19] have used the quasi-local equation of state [20, 21] to explore the stability of anisotropic stars under radial perturbations. Sharif and Kausar [22] investigated stability of expansion-free fluid in $f(R)$ gravity and found that stability of the fluid is constrained by energy density inhomogeneity, pressure anisotropy and $f(R)$ model.

The study of charged self-gravitating objects in the context of coupled Einstein-Maxwell field equations leads to the evolution of black hole [23]-[25]. The physical aspects of the electromagnetic field has a significant role in general relativity. Stettner [26] investigated the stability of a pulsating sphere with a constant surface charge. Glazer [27] generalized this result by taking an arbitrary distribution of charge and found that the Bonner's charged dust model is dynamically unstable. The stability limit for charged spheres has been proposed by many people [28]-[32] starting from the Buchdahl [33] work for neutral spheres. Recently, we have investigated the problem of dynamical instability of cylindrical and spherical systems with electromagnetic field at Newtonian and pN regimes [34].

Here we explore the role of electromagnetic field in the stability of collapsing fluid undergoing dissipation with shearing viscosity. Darmois matching conditions [35] are formulated for the continuity of interior general spherically symmetric solution to the exterior vacuum Reissner-Nordström solution. The paper is planned as: In the next section, we discuss some basic properties of the viscous fluid, Einstein-Maxwell equations and junction conditions. Sec-

tion **3** provides the perturbation scheme to form the collapse equation. In section **4**, we explore the collapse equation at Newtonian and pN regimes. Finally, we discuss our conclusion in section **5**.

2 Fluid Distribution, Field Equations and Junction Conditions

We consider a timelike three-space spherical surface, Σ, which separates the $4D$ geometries into two regions interior M^- and exterior M^+. The M^- is given by the general spherically symmetric spacetime in the comoving coordinates

$$ds^2_- = -A^2(t,r)dt^2 + B^2(t,r)dr^2 + R^2(t,r)(d\theta^2 + \sin^2\theta d\phi^2). \tag{1}$$

For the exterior metric, we take Reissner-Nordström metric describing the radiation field around a charged spherically symmetric source of the gravitational field

$$ds^2_+ = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)d\nu^2 - 2d\nu dr + r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{2}$$

where M and Q are the total mass and charge respectively. The fluid under consideration is locally dissipative in the form of shearing viscosity and the interior energy-momentum tensor of charged dissipative fluid is given by

$$T^-_{\alpha\beta} = (\rho + \hat{p} - \xi\Theta)u_\alpha u_\beta + (\hat{p} - \xi\Theta)g_{\alpha\beta} - \eta\sigma_{\alpha\beta} + \frac{1}{4\pi}\left(F^\gamma_\alpha F_{\beta\gamma} - \frac{1}{4}g_{\alpha\beta}F^{\gamma\delta}F_{\gamma\delta}\right), \tag{3}$$

where ρ, $\hat{p}$, ξ and u_α are the energy density, isotropic pressure, coefficient of bulk viscosity and four-velocity associated with the fluid, respectively, $F_{\alpha\beta}$ is the electromagnetic field tensor. We can write the above equation with $p = \hat{p} - \xi\Theta$ as

$$T^-_{\alpha\beta} = (\rho + p)u_\alpha u_\beta + pg_{\alpha\beta} - \eta\sigma_{\alpha\beta} + \frac{1}{4\pi}\left(F^\gamma_\alpha F_{\beta\gamma} - \frac{1}{4}g_{\alpha\beta}F^{\gamma\delta}F_{\gamma\delta}\right), \tag{4}$$

The four velocity in the comoving coordinates is

$$u^\alpha = A^{-1}\delta^\alpha_0, \quad u^\alpha u_\alpha = -1. \tag{5}$$

Here $\eta > 0$ is the coefficient of shearing viscosity, while the shear tensor $\sigma_{\alpha\beta}$ is defined as

$$\sigma_{\alpha\beta} = u_{(\alpha;b)} + a_{(\alpha}u_{\beta)} - \frac{1}{3}\Theta(g_{\alpha\beta} + u_\alpha u_\beta), \tag{6}$$

where $a_\alpha = u_{\alpha;\beta}u^\beta$ is the four acceleration and $\Theta = u^\alpha_{;\alpha}$ is the expansion scalar. The corresponding non-zero components turns out

$$a_1 = \frac{A'}{A}, \quad a^\alpha a_\alpha = (\frac{A'}{AB})^2, \quad \Theta = \frac{1}{A}\left(\frac{\dot{B}}{B} + 2\frac{\dot{R}}{R}\right), \tag{7}$$

where dot and prime represent differentiation with respect to t and r, respectively. The non-vanishing components for the shear tensor are

$$\sigma_{11} = 2B^2\sigma, \quad \sin^2\theta\sigma_{22} = \sigma_{33} = -R^2\sigma, \quad \sigma^{\alpha\beta}\sigma_{\alpha\beta} = 2\sigma^2, \tag{8}$$

where

$$\sigma = \frac{1}{3A}\left(\frac{\dot{R}}{R} - \frac{\dot{B}}{B}\right). \tag{9}$$

2.1 The Einstein-Maxwell Field Equations

In four-vector formalism, the Maxwell equations are

$$F_{\alpha\beta} = \phi_{\beta,\alpha} - \phi_{\alpha,\beta}, \quad F^{\alpha\beta}_{;\beta} = 4\pi J^\alpha, \tag{10}$$

where ϕ_α and J^α are the four potential and current density vector, respectively. We consider the charge to be at rest resulting no magnetic field present in this local coordinate system. Thus ϕ_α and J^α can be written as follows

$$\phi_\alpha = \Phi\delta^0_\alpha, \quad J^\alpha = \zeta u^\alpha, \tag{11}$$

where $\zeta(t,r)$ and $\Phi(t,r)$ describe the charge density and the scalar potential, respectively. From Eqs.(10) and (11), the only non-vanishing component of electromagnetic field tensor is

$$F_{10} = -F_{01} = \frac{\partial\Phi}{\partial r}. \tag{12}$$

The corresponding Maxwell equations are

$$\frac{\partial^2\Phi}{\partial r^2} - \left(\frac{A'}{A} + \frac{B'}{B} - 2\frac{R'}{R}\right)\frac{\partial\Phi}{\partial r} = 4\pi\zeta AB^2, \tag{13}$$

$$\frac{\partial^2\Phi}{\partial t\partial r} - \left(\frac{\dot{A}}{A} + \frac{\dot{B}}{B} - 2\frac{\dot{R}}{R}\right)\frac{\partial\Phi}{\partial r} = 0. \tag{14}$$

Solving the above equations, it follows that

$$\frac{\partial \Phi}{\partial r} = \frac{qAB}{R^2}, \tag{15}$$

where $q(r)$ is the total amount of charge from center to the boundary surface of the star

$$q(r) = 4\pi \int_0^r \zeta B R^2 dr. \tag{16}$$

The electric field intensity is the charge per unit surface area of the sphere

$$E(t,r) = \frac{q}{4\pi R^2}. \tag{17}$$

Using Eqs.(1) and (4), we have non-vanishing components of the Einstein-Maxwell field equations for (1) yields

$$\begin{aligned} 8\pi A^2(\rho + 2\pi E^2) &= \left(\frac{2\dot{B}}{B} + \frac{\dot{R}}{R}\right)\frac{\dot{R}}{R} - \left(\frac{A}{B}\right)^2 \\ &\times \left[\frac{2R''}{R} + \left(\frac{R'}{R}\right)^2 - \frac{2B'R'}{BR} - \left(\frac{B}{R}\right)^2\right], \end{aligned} \tag{18}$$

$$0 = -2\left(\frac{\dot{R}'}{R} - \frac{\dot{R}A'}{RA} - \frac{\dot{B}R'}{BR}\right), \tag{19}$$

$$\begin{aligned} 8\pi B^2(p + 2\eta\sigma - 2\pi E^2) &= -\left(\frac{B}{A}\right)^2\left[\frac{2\ddot{R}}{R} - \left(\frac{2\dot{A}}{A} - \frac{\dot{R}}{R}\right)\frac{\dot{R}}{R}\right] \\ &+ \left(\frac{2A'}{A} + \frac{R'}{R}\right)\frac{R'}{R} - \left(\frac{B}{R}\right)^2, \end{aligned} \tag{20}$$

$$\begin{aligned} 8\pi R^2(p - \eta\sigma + 2\pi E^2) &= 8\pi R^2(p - \eta\sigma + 2\pi E^2)\sin^{-2}\theta \\ &= -\left(\frac{R}{A}\right)^2\left[\frac{\ddot{B}}{B} + \frac{\ddot{R}}{R} - \frac{\dot{A}}{A}\left(\frac{\dot{B}}{B} + \frac{\dot{R}}{R}\right) + \frac{\dot{B}\dot{R}}{BR}\right] \\ &+ \left(\frac{R}{B}\right)^2\left[\frac{A''}{A} + \frac{R''}{R} - \frac{A'}{A}\left(\frac{B'}{B} - \frac{R'}{R}\right) - \frac{B'R'}{BR}\right]. \end{aligned} \tag{21}$$

The Misner and Sharp [29] mass function $m(t,r)$ with charge is given by

$$m(t,r)=\frac{R}{2}(1-g^{\alpha\beta}R_{,\alpha}R_{,\beta})=\frac{R}{2}\left(1+\frac{\dot{R}^2}{A^2}-\frac{R'^2}{B^2}\right)+\frac{q^2}{2R}. \tag{22}$$

The conservation equation, $(T^{-\alpha\beta})_{;\beta}=0$, yields

$$\dot{\rho}+(\rho+p)\left(\frac{\dot{B}}{B}+2\frac{\dot{R}}{R}\right)+2\eta\sigma\left(\frac{\dot{B}}{B}-\frac{\dot{R}}{R}\right)=0, \tag{23}$$

$$p'+2\eta\sigma'+(\rho+p)\frac{A'}{A}+2\eta\sigma\left(\frac{A'}{A}+3\frac{R'}{R}\right)-4\pi\frac{E}{R}(RE'+2R'E)=0. \tag{24}$$

2.2 Junction Conditions

We connect M^- and M^+ metrics by considering the Darmois junction conditions. For the smooth matching of these geometries, it is required that the boundary is continuous and smooth. Thus the continuity of first fundamental form of the metrics provides, i.e.,

$$\left(ds_-^2\right)_\Sigma=\left(ds_+^2\right)_\Sigma=\left(ds^2\right)_\Sigma, \tag{25}$$

and the continuity of second fundamental form of the extrinsic curvature gives

$$K_{ij}^+=K_{ij}^-, \quad (i,j=0,2,3). \tag{26}$$

Considering the interior and exterior spacetimes with Eq.(25), we have

$$\frac{dt}{d\tau} = A(t,r_\Sigma)^{-1}, \quad R(t,r_\Sigma)=r_\Sigma(\nu), \tag{27}$$

$$\left(\frac{d\nu}{d\tau}\right)^{-2} = \left(1-\frac{2M}{r_\Sigma}+\frac{Q^2}{r_\Sigma^2}+2\frac{dr_\Sigma}{d\nu}\right). \tag{28}$$

From Eq.(26), the non-null components of the extrinsic curvature turn out to be

$$K^{-}_{00}=-\left[\frac{A'}{AB}\right]_{\Sigma},\quad K^{-}_{22}=\left[\frac{RR'}{B}\right]_{\Sigma},\quad K^{-}_{33}=K^{-}_{22}\sin^2\theta, \tag{29}$$

$$K^{+}_{00}=\left[\left(\frac{d^2\nu}{d\tau^2}\right)\left(\frac{d\nu}{d\tau}\right)^{-1}-\left(\frac{d\nu}{d\tau}\right)\left(\frac{M}{r^2}-\frac{Q^2}{r^3}\right)\right]_{\Sigma}, \tag{30}$$

$$K^{+}_{22}=\left[\left(\frac{d\nu}{d\tau}\right)\left(1-\frac{2M}{r}-\frac{Q^2}{r^2}\right)r+\left(\frac{dr}{d\tau}\right)r\right]_{\Sigma}, \tag{31}$$

$$K^{+}_{33}=K^{+}_{22}\sin^2\theta. \tag{32}$$

Using Eqs.(27)-(31) and the field equations, we obtain

$$m\overset{\Sigma}{=}M,\quad p+2\eta\sigma\overset{\Sigma}{=}0. \tag{33}$$

where $q(r)=Q$ has been used. The above equation shows that across the boundary, Σ, the masses of interior and exterior spacetimes are matched as well as momentum flux is conserved.

3 Perturbation Scheme and Collapse Equation

Here we construct the collapse equation. For this purpose, we perturb the field equations, dynamical equations and the mass function upto first order in ε by using the perturbation scheme [22, 34]. We assume that initially all the physical functions and the metric coefficients depend on r, i.e., the fluid is unperturbed. Afterwards, all these quantities depend on time coordinate, which are given by

$$A(t,r) = A_0(r)+\varepsilon T(t)a(r), \tag{34}$$

$$B(t,r) = B_0(r)+\varepsilon T(t)b(r), \tag{35}$$

$$R(t,r) = rB(t,r)[1+\varepsilon T(t)\bar{c}(r)], \tag{36}$$

$$E(t,r) = E_0(r)+\varepsilon T(t)e(r), \tag{37}$$

$$\rho(t,r) = \rho_0(r)+\varepsilon\bar{\rho}(t,r), \tag{38}$$

$$p(t,r) = p_0(r)+\varepsilon\bar{p}(t,r), \quad (39)$$
$$\sigma(t,r) = \varepsilon\bar{\sigma}(t,r), \quad (40)$$
$$m(t,r) = m_0(r)+\varepsilon\bar{m}(t,r), \quad (41)$$

where $0<\varepsilon\ll 1$. For $\bar{c}=0$, we have shearfree metric. Using this scheme, the static configuration of Eqs.(18)-(21) are written as

$$8\pi\left(\rho_0+2\pi E_0^2\right) = -\frac{1}{B_0^2}\left[2\frac{B_0''}{B_0}-\left(\frac{B_0'}{B_0}\right)^2+\frac{4}{r}\frac{B_0'}{B_0}\right], \quad (42)$$
$$8\pi\left(p_0-2\pi E_0^2\right) = \frac{1}{B_0^2}\left[\left(\frac{B_0'}{B_0}\right)^2+\frac{2}{r}\left(\frac{A_0'}{A_0}+\frac{B_0'}{B_0}\right)+2\frac{A_0'}{A_0}\frac{B_0'}{B_0}\right], \quad (43)$$
$$8\pi\left(p_0+2\pi E_0^2\right) = \frac{1}{B_0^2}\left[\frac{A_0''}{A_0}+\frac{B_0''}{B_0}+\frac{1}{r}\left(\frac{A_0'}{A_0}+\frac{B_0'}{B_0}\right)-\left(\frac{B_0'}{B_0}\right)^2\right]. \quad (44)$$

The corresponding perturbed quantities upto first order in ε with Eqs.(34)-(40) become

$$8\pi\bar{\rho}+32\pi^2E_0Te = -\frac{T}{B_0^3}\left\{b\left(\frac{B_0'}{B_0}\right)^2-2b'\left(\frac{B_0'}{B_0}-\frac{2}{r}\right)+2b''+2B_0 \times \left[\bar{c}''+\bar{c}'\left(2\frac{B_0'}{B_0}+\frac{3}{r}\right)+\left(\frac{\bar{c}}{r^2}\right)\right]\right\} - 24\pi\frac{Tb}{B_0}\left(\rho_0+2\pi E_0^2\right), \quad (45)$$
$$0 = 2\left[\left(\frac{b}{A_0B_0}\right)'+\left(\frac{\bar{c}}{A_0}\right)'+\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{\bar{c}}{A_0}\right)\right]\dot{T}, \quad (46)$$
$$8\pi(\bar{p}+2\eta\bar{\sigma})-32\pi^2E_0Te = -\frac{2b}{A_0^2B_0}\ddot{T}+\frac{2T}{B_0^2}\left[\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{a}{B_0}\right)' + \left(\frac{b}{B_0}+\bar{c}\right)'\left(\frac{A_0'}{A_0}+\frac{1}{r}+\frac{B_0'}{B_0}\right)+\frac{\bar{c}}{r^2} - \bar{c}\left(\frac{B_0}{A_0}\right)^2\frac{\ddot{T}}{T}\right]-16\pi\frac{Tb}{B_0}\left(p_0-2\pi E_0^2\right), \quad (47)$$

$$
\begin{aligned}
8\pi(\bar{p}-\eta\bar{\sigma})+32\pi^2E_0Te &= -\frac{\ddot{T}}{A_0^2}\left[2\frac{b}{B_0}+\bar{c}\right]+\frac{T}{B_0^2}\left[\frac{a''}{A_0}+\frac{b''}{B_0}+\bar{c}''\right.\\
&+ \frac{A_0''}{A_0}\left(\frac{a}{A_0}-\frac{b}{B_0}\right)+\frac{b'}{B_0}\left(\frac{1}{r}-\frac{B_0'}{B_0}\right)\\
&+ \frac{B_0'}{B_0}\left(\bar{c}-\frac{b}{B_0}\right)'+\frac{A_0'}{A_0}\left(\frac{b}{rB_0}+\bar{c}'\right)\\
&+ \left.\frac{1}{r}\left(\frac{a}{A_0}+2\bar{c}\right)'\right]-24\pi\frac{Tb}{B_0}\left(p_0+2\pi E_0^2\right). \quad (48)
\end{aligned}
$$

The static configuration for the dynamical equation (23) is trivially satisfied, while Eq.(24) implies that

$$
p_0'+(\rho_0+p_0)\frac{A_0'}{A_0}-4\pi E_0\left[E_0'+2E_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\right]=0, \quad (49)
$$

which yields

$$
\frac{A_0'}{A_0} = -\frac{1}{\rho_0+p_0}\left\{p_0'-4\pi E_0\left[E_0'+2E_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\right]\right\}. \quad (50)
$$

The perturbed part of Eq.(23) becomes

$$
\dot{\bar{\rho}}+(\rho_0+p_{r0})\left(3\frac{b}{B_0}+2\bar{c}\right)\dot{T}=0, \quad (51)
$$

which on integration leads to

$$
\bar{\rho}=-(\rho_0+p_0)\left(3\frac{b}{B_0}+2\bar{c}\right)T. \quad (52)
$$

For the second dynamical equation, we have

$$
\begin{aligned}
&\bar{p}'+2\eta\bar{\sigma}'+(\bar{\rho}+\bar{p})\frac{A_0'}{A_0}+(\rho_0+p_{r0})T\left(\frac{a}{A_0}\right)'\\
+ \;& 2\eta\sigma\left(\frac{A_0'}{A_0}+\frac{3}{r}+\frac{3B_0'}{B_0}\right)-4\pi T\left[(eE_0)'+4eE_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\right.\\
+ \;& \left.2E_0^2\left(\frac{b}{B_0}\right)'+2E_0^2\bar{c}'\right]=0. \quad (53)
\end{aligned}
$$

The unperturbed and perturbed configurations of electromagnetic field are given by

$$E_0(r)=\frac{q}{r^2B_0^2},\quad e(r)=-\left(\frac{2b}{B_0}+\bar{c}\right)E_0. \tag{54}$$

This shows that $e(r)$ depends upon the static configuration of electromagnetic field. Similarly, we can write the static and perturbed configuration for Eq.(22) as

$$\begin{aligned} m_0 &= -\frac{rB_0}{2}\left(2r\frac{B_0'}{B_0}+\left(r\frac{B_0'}{B_0}\right)^2\right)+8\pi^2r^3E_0^2B_0^3, \qquad (55)\\ \bar{m} &= -\left[r^2b'+r^3\frac{B_0'^2}{2B_0}\left(2\frac{b'}{B_0'}-\frac{b}{B_0}\right)\right]T-\bar{c}'r^3B_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)T\\ &- \bar{c}r^2B_0\left[\frac{3}{2}r\left(\frac{B_0'}{B_0}\right)^2+3\frac{B_0'}{B_0}+\frac{1}{r}\right]T+24\pi^2r^3B^2E_0^2\left(b+B_0\bar{c}\right)T. \qquad (56)\end{aligned}$$

From Eq.(55), it follows that

$$\frac{B_0'}{B_0}=-\frac{1}{r}+\frac{1}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2} \tag{57}$$

The perturbed configuration for the shear scalar is written as

$$\bar{\sigma}=\frac{1}{3A_0}\bar{c}\dot{T}. \tag{58}$$

The junction condition (33) with Eqs.(39) and (40) leads to

$$p_0\overset{\Sigma}{=}0,\quad \bar{p}+2\eta\bar{\sigma}\overset{\Sigma}{=}0. \tag{59}$$

Substituting the above relations in Eq.(47), we get temporal equation

$$\ddot{T}(t)-\psi(r)T(t)\overset{\Sigma}{=}0, \tag{60}$$

where

$$\begin{aligned}\psi(r) &\overset{\Sigma}{=} \left(\frac{A_0}{B_0}\right)^2\left(\frac{b}{B_0}+\bar{c}\right)^{-1}\left[\left(\frac{A_0'}{A_0}+\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{b}{B_0}+\bar{c}\right)'\right.\\ &+ \left.\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{a}{A_0}\right)'+\frac{\bar{c}}{r^2}+16\pi^2E_0B_0(eB_0+bE_0)\right]. \qquad (61)\end{aligned}$$

The general solution of Eq.(60) yields

$$T(t) = c_1 \exp(\sqrt{\psi_\Sigma} t) + c_2 \exp(-\sqrt{\psi_\Sigma} t), \tag{62}$$

where c_1 and c_2 are arbitrary constants. This provides two independent solutions. Here we take ψ_Σ to be positive for physically meaningful result, i.e., when the system is in static position, it starts collapsing at $t = -\infty$ when $T(-\infty) = 0$ and goes on collapsing diminishing its areal radius with the increase of t. The corresponding solution is found for $c_2 = 0$, also, we choose $c_1 = -1$, hence

$$T(t) = -\exp(\sqrt{\psi_\Sigma} t). \tag{63}$$

Next, we are interested to find instability range of the collapsing fluid in terms of adiabatic index Γ. Chan et al. [16] found a relationship between $\bar{p}$ and $\bar{\rho}$ with an equation of state of Harrison-Wheeler type [36] for the static configuration as

$$\bar{p} = \Gamma \frac{p_0}{\rho_0 + p_0} \bar{\rho}, \tag{64}$$

where Γ describes the change in pressure for a given change in density (taken to be constant in the whole fluid distribution). The above equation with Eq.(52) leads to

$$\bar{p} = -p_0 \Gamma \left(\frac{3b}{B_0} + 2\bar{c} \right) T. \tag{65}$$

Using the above equation and (59) in Eq.(47), it follows that

$$\begin{aligned} \left(\frac{a}{A_0} \right)' &= B_0 \left(\frac{1}{r} + \frac{B_0'}{B_0} \right)^{-1} \left[-4\pi \left(\frac{3b}{B_0} + 2\bar{c} \right) \Gamma p_0 B_0 + 8\pi p_0 b - \frac{\bar{c}}{r^2 B_0} \right. \\ &- 16\pi^2 E_0 (eB_0 + bE_0) - \frac{1}{B_0} \left(\frac{A_0'}{A_0} + \frac{1}{r} + \frac{B_0'}{B_0} \right) \left(\frac{b}{B_0} + \bar{c} \right)' \\ &+ \left. \frac{B_0}{A_0^2} \left(\frac{b}{B_0} + \bar{c} \right) \frac{\ddot{T}}{T} + 8\pi\eta B_0 \frac{\bar{\sigma}}{T} \right]. \end{aligned} \tag{66}$$

We can write the collapse equation by substituting Eqs.(52), (58), (65) and (66) in the perturbed configuration of the second dynamical equation as follows

$$- \left[p_0 \Gamma \left(\frac{3b}{B_0} + 2\bar{c} \right) \right]' T + \frac{2}{3A_0} \left(\eta \bar{c}' + \eta \bar{c} \frac{A_0'}{A_0} \right) \dot{T} - \left(\frac{3b}{B_0} + 2\bar{c} \right) T$$

$$\times \quad (\rho_0+p_0(1+\Gamma))\,\frac{A_0'}{A_0}+(\rho_0+p_0)TB_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)^{-1}\left[\frac{B_0}{A_0^2}\left(\frac{b}{B_0}+\bar{c}\right)\frac{\ddot{T}}{T}\right.$$
$$+ \quad 8\pi p_0 b+8\pi\eta B_0\frac{\bar{c}\dot{T}}{3A_0T}-16\pi^2E_0(eB_0+bE_0)-4\pi\left(\frac{3b}{B_0}+2\bar{c}\right)\Gamma p_0B_0$$
$$\left.- \quad \frac{1}{B_0}\left(\frac{A_0'}{A_0}+\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{b}{B_0}+\bar{c}\right)'-\frac{\bar{c}}{r^2B_0}\right]+\frac{2}{3A_0}\left(\frac{A_0'}{A_0}+\frac{1}{r}+\frac{B_0'}{B_0}\right)$$
$$\times \quad \eta\bar{c}\dot{T}-T\left[2E_0^2\left(\frac{b}{B_0}\right)'+(eE_0)'+2E_0^2\bar{c}'+4\pi E_0\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\right]=0. \tag{67}$$

4 Dynamical Instability of Charged Viscous Perturbation

This section deals with the dynamical instability of charged viscous fluid in the frame work of Newtonian and pN regimes. We see from Eq.(36) that the effects of shear explicitly appear only on metric function $R(t,r)$. Using this fact, we can split Eq.(46) in two types of terms one with shearing viscosity and other without it. Thus, the first term as well as the second and third terms of Eq.(46) are taken identically equal to zero

$$\left(\frac{b}{A_0B_0}\right)'=0, \quad \left(\frac{\bar{c}}{A_0}\right)'+\left(\frac{1}{r}+\frac{B_0'}{B_0}\right)\left(\frac{\bar{c}}{A_0}\right)=0. \tag{68}$$

The first of the above equation provides

$$b=A_0B_0, \tag{69}$$

while the second leads to

$$\bar{c}' \quad = \quad -\bar{c}\left\{\left[\frac{p_0'}{\rho_0+p_0}+\frac{4\pi E_0}{\rho_0+p_0}\left(E_0'+\frac{2E_0}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right]\right.$$
$$\left. + \quad \frac{1}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right\}. \tag{70}$$

Inserting Eqs.(50), (57), (63), (69) and (70) in the collapse equation, we have

$$
\begin{aligned}
& -(3A_0+2\bar{c})\Gamma p_0' + \Gamma p_0\left\{A_0\left[\frac{3p_0'}{\rho_0+p_0} - \frac{12\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\right.\right.\right. \\
+ & \left.\left.\left.\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right] + \frac{2p_0'\bar{c}}{\rho_0+p_0} + \frac{2\bar{c}}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2} \\
- & \left.\frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right\} - (\rho_0+p_0+\Gamma p_0) \\
\times & (3A_0+2\bar{c})\left[\frac{p_0'}{\rho_0+p_0} - \frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right] \\
+ & (\rho_0+p_0)B_0\left(\frac{1}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)^{-1}\left\{\frac{8\pi}{3}\frac{B_0}{A_0}\eta\bar{c}\sqrt{\psi_\Sigma}\right. \\
- & \left.16\pi^2E_0(eB_0+bE_0) + 8\pi p_0A_0B_0 - 12\pi\Gamma p_0A_0B_0 + \left(\frac{B_0}{A_0}\right)\psi_\Sigma\right. \\
+ & \left(\frac{A_0}{B_0}\right)\left[\frac{1}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2} - \frac{p_0'}{\rho_0+p_0} + \frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\right.\right. \\
\times & \left.\left.\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right]\left[\frac{p_0'}{\rho_0+p_0} - \frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\right.\right. \\
\times & \left.\left.\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right] + \frac{\bar{c}}{B_0}\left[\frac{p_0'}{\rho_0+p_0} - \frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\right.\right. \\
\times & \left.\left.\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right]^2 - \frac{\bar{c}}{B_0}\left(\frac{2m_0}{r^3B_0} - 16\pi^2E_0^2B_0^2 + 8\pi B_0^2p_0\Gamma\right. \\
- & \left.\left.\left(\frac{B_0}{A_0}\right)^2\psi_\Sigma\right)\right\} + \frac{2}{3A_0r}\eta\bar{c}\sqrt{\psi_\Sigma}\left[2\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2} - \frac{p_0'}{\rho_0+p_0}\right. \\
+ & \left.\frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right] - \{(eE_0)' + 2(A_0+\bar{c}) \\
\times & \left[-\frac{p_0'}{\rho_0+p_0} + \frac{4\pi E_0}{\rho_0+p_0}\left(E_0' + \frac{2E_0}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right)\right]E_0^2 \\
+ & \left.\frac{(4\pi E_0+2E_0^2\bar{c})}{r}\sqrt{1-\frac{2m_0}{rB_0}+16\pi^2r^2E_0^2B_0^2}\right\} = 0. \qquad (71)
\end{aligned}
$$

4.1 Newtonian Limit

Now we explore the instability range by applying the Newtonian limit, i.e., $A_0 = 1,\ B_0 = 1$ and $\rho_0 \gg p_0$ and discarding the terms of the order $\frac{m_0}{r}$ in Eq.(71). It follows that

$$-3p_0'\Gamma + 4p_0' + \frac{(1+8\pi^2r^2E_0^2)}{r}\left(\frac{4}{3}\eta\bar{c}\sqrt{\psi_\Sigma} - 32\pi E_0^2 - 4\pi E_0\right)$$
$$- \ 16\pi E_0E_0' + (1-16\pi^2r^2E_0^2)r\rho_0\psi_\Sigma = 0. \tag{72}$$

We see from Eq.(58) that the perturbed configuration of the shear scalar depends on the velocity gradients. Also, it is known that the velocity of the particles for a self-gravitating star increases towards the center. This shows that $\bar{\sigma}$ has negative value inside the body and zero on the boundary surface. It is worth noticing that for a collapsing body $\dot{T} < 0$ as $t \to -\infty$, thus Eq.(58) implies that $\bar{c} > 0$. Considering the physical requirement, i.e., $p_0' < 0$ and neglecting the terms like $\frac{\rho_0}{p_0}$ being relativistic terms, we have instability condition (independent of linear perturbation functions) for the charged viscous fluid as

$$\Gamma < \frac{4}{3} - \frac{4}{9}\eta|\bar{c}|\frac{\sqrt{\psi_\Sigma}}{|p_0'|r} + \frac{16\pi E_0E_0'}{3|p_0'|} + \frac{1+8\pi^2r^2E_0^2}{3|p_0'|r}(32\pi E_0^2 + 4\pi E_0). \tag{73}$$

Here the critical value $\frac{4}{3}$ corresponds to the spherical geometry and Newtonian gravity. In fact, 4 in the numerator corresponding to the weight of the envelop in Newtonian mechanics varying as r^{-2} which is distributed over the surface of sphere yielding another r^{-2}. The denominator 3 corresponds to the volume of the sphere r^3. We note from the above equation that the electromagnetic field diminishes the impact of shearing viscosity on the dynamical instability and makes the system unstable at Newtonian approximation. We retain the Newtonian classical result, i.e., $\Gamma < \frac{4}{3}$, for the case of shearfree or when fluid is not charged viscous.

4.2 Post-Newtonian Limit

In view of pN limit, we consider $A_0 = 1-\frac{m_0}{r},\ B_0 = 1+\frac{m_0}{r}$ and the relativistic corrections of the order $\frac{m_0}{r}$ in Eq.(71) so that we have

$$- \ (3+2\bar{c})p_0'\Gamma + (4+2\bar{c})p_0' + \frac{(1+8\pi^2r^2E_0^2)}{r}\left(\frac{4}{3}\eta\bar{c}\sqrt{\psi_\Sigma} + 4\pi E_0\right.$$
$$+ \ \left.2E_0^2\bar{c} - 8\pi E_0^2(2+2\bar{c})\right) + (1-16\pi^2r^2E_0^2)r\rho_0[8\pi p_0 + \psi_\Sigma$$

$$
\begin{aligned}
&- \quad 16\pi^2 E_0(e+E_0)+\frac{8\pi}{3}\eta\bar{c}\sqrt{\psi_\Sigma}]-(4+2\bar{c})4\pi E_0 E_0'-(eE_0)' \\
&+ \quad 16\pi^2 r\bar{c}\rho_0 E_0^2+\rho_0 r\bar{c}\psi_\Sigma(1-8\pi^2 r^2 E_0^2)=0. \qquad (74)
\end{aligned}
$$

Here we apply the same procedure as in the Newtonian limit and ignoring terms with higher order $\frac{m_0}{r}$, we get the instability range at pN limit as follows

$$
\begin{aligned}
\Gamma < \frac{4}{3}-\frac{1}{3|p_0'|}&\left[\left(\frac{4}{3r}\eta|\bar{c}|\sqrt{\psi_\Sigma}+4\pi E_0+2E_0^2|\bar{c}|\right)(1+8\pi^2 r^2 E_0^2)\right. \\
+ \quad & 16\pi^2 E_0^2 r\rho_0(|\bar{c}|+16\pi^2 E_0 r^2(e+E_0))+2|\bar{c}|+\rho_0\psi_\Sigma r(1+|\bar{c}|) \\
+ \quad & \left.\frac{8\pi}{3}\eta\rho_0 r|\bar{c}|\sqrt{\psi_\Sigma}\right]+\frac{16\pi^2 r E_0\rho_0}{3|p_0'|}\left[\frac{8\pi}{3}\eta|\bar{c}|r^2 E_0\sqrt{\psi_\Sigma}+e+E_0\right. \\
+ \quad & \left.r^2 E_0\psi_\Sigma(1+\frac{|\bar{c}|}{2})\right]+\frac{1}{3|p_0'|}\left[8\pi\rho_0 p_0 r+4\pi E_0 E_0'(4+2|\bar{c}|)+(eE_0)'\right. \\
+ \quad & \left.16\pi E_0^2(1+|\bar{c}|)\frac{(1+8\pi^2 r^2 E_0^2)}{r}\right]. \qquad (75)
\end{aligned}
$$

This equation shows that the positive terms diminish the relativistic effects of negative terms occurring from the shearing viscosity and electromagnetic field at pN approximation, making the fluid more unstable. It is mentioned here that instability condition depends upon the static configuration of the system, as perturbed variable $e(r)$ and $b(r)$ depend on static configuration given in Eqs.(54) and (69) respectively.

5 Conclusion

We have investigated the role of electromagnetic field on the instability conditions (73) and (75) at Newtonian and pN approximations. Pinheiro and Chan [37] found that the collapsing stars with huge amount of charge ($Q \sim 5.408 \times 10^{20}$ Coulomb) leads to the Reissner-Nordström black hole. They concluded that the models with charge to mass ratio $\frac{Q}{m_0} \leq 0.631$ would form a black hole. Ernesto and Simeone [38] examined the stability of charged thin shell and found that Reissner-Nordström geometry for different values of charge either have an inner and outer event horizon or a naked singularity. This shows the relevance of charge on the evolution and structure formation of astrophysical objects.

In general, the behavior of electromagnetic field is always positive being the Coulomb's repulsion force. Sharif and Abbas [1] explored that due to

the weak nature of the electromagnetic field, the end state of collapsing cylinder results a charged black string. We see from Eq.(73) that shearing viscosity boost the stability of the fluid which is the consequence of the fact that the collapse with shear proceeds faster than without shear. Whereas the electromagnetic field being a positive quantity diminishes the effects of shearing viscosity and makes the fluid unstable at Newtonian approximation. This corresponds to the fact that charge delays the event horizon formation or even halts the complete contraction of the star [10]. Also, the electromagnetic field has the same impact for Eq.(75) at the pN approximation. It is worth mentioning here that our results for electromagnetic field are consistent with the results obtained in [34].

Finally, we would like to mention that we have made all the discussion on the hypersurface, where the areal radius is constant. The solution of the temporal equation (60) includes oscillating and non-oscillating functions correspond to stable and unstable systems. For the sake of instability conditions, we have confined our interest in the non-oscillating ones. We have investigated the role of physical quantities in the onset of dynamical instability of fluid during the collapse, hence the instability conditions contain those terms which have radial dependence.

Acknowledgments

We would like to thank the Higher Education Commission, Islamabad, Pakistan, for its financial support through the *Indigenous Ph.D. 5000 Fellowship Program Batch-VII*. One of us (MA) would like to thank University of Education, Lahore for the study leave.

References

[1] Sharif, M. and Abbas, G.: Astrophys. Space Sci. **327**(2010)285; J. Phys. Soc. Jpn. **80**(2011)104002; Chinese Phys. B **22**(2013)030401.

[2] Eddington, A.S.: *Internal Constitution of the Stars* (Cambrigde University Press, 1926).

[3] Glendenning, N.: *Compact Stars* (Springer, 2000).

[4] Rosseland, S.: Mon. Not. R. Astron. Soc. **84**(1924)720.

[5] de la Cruz, V. and Israel, W.: Nuovo Cimento A **51** (1967)744.

[6] Bekenstein, J.: Phys. Rev. D **4**(1971)2185.

[7] Olson, E. and Bailyn, M.: Phys. Rev. D **13**(1976)2204.

[8] Mashhoon, B. and Partovi, M.: Phys. Rev. D **20**(1979)2455.

[9] Zhang, J.L., Chau, W.Y. and Deng, T.Y.: Astrophys. Space Sci. **88**(1982)81.

[10] Ghezzi, C.: Phys. Rev. D **72**(2005)104017.

[11] Barreto, W, Rodrguez, B., Rosales, L and Serrano, O.: Gen. Relativ. Gravit. **39**(2007)537.

[12] Chandrasekhar, S.: Astrophys. J. **140**(1964)417.

[13] Herrera, L., Santos, N.O. and Le Denmat, G.: Mon. Not. R. Astron. Soc. **237**(1989)257.

[14] Chan, R., Kichenassamy, S., Le Denmat, G. and Santos, N.O.: Mon. Not. R. Astron. Soc. **239**(1989)91.

[15] Chan, R., Herrera, L. and Santos, N.O.: Mon. Not. R. Astron. Soc. **265**(1993)533.

[16] Chan, R., Herrera, L. and Santos, N.O.: Mon. Not. R. Astron. Soc. **267**(1994)637.

[17] Herrera, L., Santos, N.O. and Le Denmat, G.: Gen. Relativ. Gravit. **44**(2012)1143.

[18] Chan, R.: Mon. Not. R. Astron. Soc. **316**(2000)588.

[19] Horvat, D., Ilijic, S. and Marunovic, A.: Class. Quantum Grav. **28**(2011)25009.

[20] Hernandez, H., Nunez, L.A. and Percoco, U.: Class. Quantum Grav. **16**(1999)871.

[21] Hernandez, H., Nunez, L.A.: Can. J. Phys. **82**(2004)29.

[22] Sharif, M. and Kausar, H.R.: JCAP **07**(2011)022; Astrophys. Space Sci. **337**(2012)805.

[23] De Felice, F., Yu, Y. and Fang, Z.: Mon. Not. R. Astron. Soc. **277**(1995)L17.

[24] De Felice, F., Siming, L. and Yunqiang, Y.: Class. Quantum Grav. **16**(1999)2669.

[25] De Felice, F., Siming, L. and Yunqiang, Y.: Phys. Rev. D **68**(2003)084004.

[26] Stettner, R.: Ann. Phys. **80**(1973)212.

[27] Glazer, I.: Ann. Phys. **101**(1976)594.

[28] Mak, M., Dobson, P. and Harko, T.: Europhys. Lett. **55**(2001)310.

[29] Misner, C.W. and Sharp, D.: Phys. Rev. **136**(1964)B571.

[30] Giuliani, A. and Rothman, T.: Gen. Relativ. Gravit. **40**(2008)1427.

[31] Andreasson, H.: Commun. Math. Phys. **288**(2009)715.

[32] Bohmer, C. and Harko, T.: Gen. Relativ. Gravit. **39**(2007)757.

[33] Buchdahl, H.: Phys. Rev. **116**(1959)1027.

[34] Sharif, M. and Azam, M.: JCAP **02**(2012)043; Gen. Relativ. Gravit. **44**(2012)1181.

[35] Darmois, G.: Memorial des Sciences Mathematiques (Gautheir-Villars, 1927) Fasc. 25.

[36] Harrison, B.K., Thorne, K.S., Wakano, M. and Wheeler, J. A.: *Gravitation Theory and Garvitational Collapse* (University of Chicago Press, 1965)

[37] Pinheiro, G. and Chan, R.: Gen. Relativ. Gravit. **45**(2013)243.

[38] Ernesto, F.E. and Simeone, C.: Phys. Rev. D **83**(2011)104009.